

**Effective against
SARS-CoV-2²
in 1 minute**

STRIKE FORCE™

One-Step RTU Disinfectant

For Hospitals, Restaurants, Institutional and Industrial Use

This product is an EPA registered, ready-to-use disinfectant-cleaner-virucide-fungicide-mildewstat-deodorizer for use on hard, nonporous surfaces in many types of facilities:

- **Medical:** hospitals, nursing homes, senior care and residential facilities, doctor's offices, dental offices, surgical centers, urgent care facilities
- **Educational:** schools, universities, daycare centers, kindergartens, playgrounds
- **Foodservice and Hospitality:** hotels, convention centers, restaurants, bars, cafeterias, catering operations, food and beverage processing plants
- **Retail:** retail stores, grocery stores, gas stations, department stores, salons, barber shops, banks, office buildings
- **Fitness:** health clubs, gyms, yoga facilities, martial arts operations, spas, hockey rinks
- **Government:** jails, correctional facilities, government buildings
- **Transportation:** car rental, airplanes, airports, buses, public transportation, trains, boats, cruise ships
- **Animal Care:** veterinary practices, zoos, kennels, pet shops

Effective against a broad spectrum of listed bacteria, viruses, fungi and animal viruses:

Bacteria: *Acinetobacter baumannii*, *Campylobacter jejuni*, *Enterobacter aerogenes*, *Enterococcus faecalis*, *Enterococcus faecalis* - Vancomycin Resistant (VRE), *Escherichia coli*, *ESBL Escherichia coli*, *Klebsiella pneumoniae*, *Legionella pneumophila*, *Listeria monocytogenes*, *Pseudomonas aeruginosa*, *Salmonella enterica*, *Salmonella typhi*, *Serratia marcescens*, *Shigella dysenteriae*, *Staphylococcus aureus*, *Staphylococcus aureus* - Community Associated Methicillin Resistant (CA-MRSA), *Staphylococcus aureus* - Methicillin-Resistant (MRSA), *Staphylococcus aureus* - Vancomycin Intermediate Resistant (VISA), *Streptococcus pyogenes*, *Vibrio cholerae*.

Viruses: *Hepatitis B Virus (HBV), *Hepatitis C Virus (HCV), *Herpes Simplex Virus Type 1, *Herpes Simplex Virus Type 2, *HIV-1 (AIDS Virus), *Human Coronavirus, *Influenza Virus Type A/Brazil, *Influenza Virus Type A/Hong Kong, *2013 Influenza A Virus (H7N9), *Respiratory Syncytial Virus, *SARS Associated Coronavirus (SARS), *SARS-Related Coronavirus 2 (cause of COVID-19), *Vaccinia, *Norwalk Virus (Norovirus), *Rotavirus.

Fungi: *Aspergillus niger*, *Aspergillus fumigatus*, *Trichophyton mentagrophytes* (the Athlete's Foot Fungus)

Animal Viruses: Canine Distemper Virus, Feline Calicivirus, Murine Norovirus, Newcastle's Disease Virus, Pseudorabies Virus.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected.

DISINFECTION/VIRUCIDAL*/FUNGICIDAL/MOLD & MILDEW CONTROL DIRECTIONS: Apply this product undiluted to hard, non-porous surfaces, thoroughly wetting the surfaces with a cloth, mop, sponge, sprayer or by immersion. Treated surfaces must remain visibly wet for 10 minutes. **For SARS-CoV-2**, treated surfaces must remain visibly wet for 1 minute.

For Influenza Virus Type A and Human Coronavirus, treated surfaces must remain visibly wet for 1 minute. Wipe dry with a cloth, sponge or mop or allow to air dry. For visibly soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse spray device. Spray 6-8 inches from surface. Avoid breathing spray. Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes.

For Fungicidal Activity: This product is an effective one-step fungicide against *Trichophyton mentagrophytes* when used on hard, non-porous surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. Allow surfaces to remain visibly wet for 10 minutes.

For Mold and Mildew: Repeat treatment every 7 days or more often if new growth appears.

VETERINARY PRACTICE/ANIMAL CARE/ZOOS/PET SHOPS/KENNELS DISINFECTION/VIRUCIDAL* DIRECTIONS: For cleaning and disinfecting hard, non-porous surfaces: equipment used for feeding or watering animals, utensils, instruments, cages, kennels, stables, catteries, pens, stalls and etc. Consult product label for cleaning steps required before using this product. After cleaning, saturate surfaces with this product for a period of 10 minutes. Wipe or allow to air dry. See product label for additional instructions.

TO CLEAN AND DEODORIZE: Spray directly on surface to be cleaned. Wipe or allow to air dry.

TO SANITIZE SOFT SURFACES: Hold bottle 6" - 8" from surface and spray until fabric is wet. Fabric must remain visibly wet for 5 minutes. Allow fabric to air dry. Visibly soiled areas must be cleaned prior to sanitizing. See product label.

Safety Reminder

Consult product label and Safety Data Sheet (SDS) before use. SDS's may be downloaded from www.crown-chem.com.



Technical Specifications

Appearance	Blue Liquid
Odor	Ocean Fresh
pH (full strength, 22°C)	<11.0
Foaming	Moderate
Flash Point	None
Density (lbs/gal, 22°C)	8.33 ±0.1
Stability Range	40-100°F
Free Alkalinity (as Na2O)	Low
Solubility (in H2O, 22°C)	100%
Phosphate Content (as P)	None

Product Number

33006
33041
33005
33030
33055
330TB

Packaging

6x1 qt. case*
4x1 gal. case
5 gal. pail
30 gal. drum
55 gal. drum
275 gal. tote

*One sprayer included per case